



INTRODUCTION

Agricultural activities are receiving an increasing amount of attention for allegedly contributing excess nutrients to our country's water resources. Nutrient management planning, together with proper land application, can reduce adverse impacts. Potential impacts are eliminated as growers learn to implement proper soils and litter management procedures.

Nitrogen

Most of the nitrogen found in poultry manure or litter is in the form of organic nitrogen. A smaller amount of the nitrogen in the manure is ammonium. Organic nitrogen can be mineralized or converted by soil bacteria into inorganic nitrogen. In this form, nitrogen is readily available for plant uptake. Excessive organic and ammonium forms of nitrogen can be transformed into nitrate nitrogen. High levels of nitrate can be harmful to human health. Excess nitrogen can be removed from application sites by surface runoff and leaching, and then flow into surface and ground water sources.

Phosphorous



Poultry manure is also composed of relatively large amounts of phosphorus. Phosphorus is essential for plant and animal nutrition. In the soil, phosphorus is made-up of inorganic and organic forms. Both forms interact to release water soluble phosphorous which can then be used to meet the nutritional requirements of the plants and vegetation.

Phosphorous laden soils can be eroded by rainfall and the particles can then be transported into surface water sources. Excessive phosphorus in surface waters can cause excessive plant and algal growth. Excessive algal growth can contribute to fish kills by depleting the dissolved oxygen content of the water.

Producers must implement proper land application management practices for manures with high phosphorous concentrations. Adopting conservation practices which reduce soil erosion will lower the potential for phosphorus being introduced into surface waters. Manure application rates based on nitrogen plant requirements can lead to phosphorous buildup in soils. The practice of rotating crops and application sites will help remove excess phosphorus. Maintaining soil pH between 6.0 and 7.0, maximizes plant phosphorus uptake, thereby reducing accumulations.

Producers should also implement an annual soil sampling program for application fields to determine nutrient concentrations and to help calculate application rates. Further applications should not be made to soils containing excessive phosphorus amounts.





Potassium

Potassium is another important nutrient contained in poultry manure. Potassium is rather mobile in the soil, but is available to supply plant nutritional requirements. Excessive potassium, though, can cause adverse plant health impacts. Proper soil monitoring is important for ensuring plant production.

Heavy Metal and Trace Elements

Poultry manure can contain trace amounts of copper, selenium, nickel, lead, and zinc. Over application of manure with high concentrations of these minerals can exceed the soil's adsorptive capacity. In addition to harming plant health, over application can increase the potential for water impacts.

Usage of Manure Nutrients

The nutrients in poultry litter are an excellent soil conditioner and fertilizer. Poultry growers must maximize the benefits of these nutrients while minimizing impacts to ground and surface water sources. Soil and litter sampling results allow producers to monitor their nutrients to meet plant needs and preserve water quality.







Best Management Practices Dry Litter Disposal & Dead Bird Disposal

Tyson has a long-standing commitment to protecting the environment wherever we have operations. This applies to our processing plants, hatcheries, feed mills, and all other aspects of live production. Due primarily to increasing concern for safe drinking water, Federal and State governments are under considerable pressure to regulate and/or mandate restrictions on the use of animal wastes. Unless both Tyson and our Producers act to address these concerns, the use of poultry litter could be severely restricted in many locations. Dead bird disposal methods have also been addressed for the same reasons.

The following requirements are necessary to optimize our environmental stewardship. Any Federal, State or local regulations in these areas take precedence and must be complied with.

Dry Litter Disposal



Regulations

Federal, State, and County regulations are becoming more prevalent in dry manure management. Where these regulations are in place, it is our responsibility to follow them completely. Tyson feels a responsibility to communicate and educate all Producers to be good stewards of the environment by the use of Best Management Practices.

Regardless of regulatory requirements, it is Tyson's expectation that we minimize the environmental impact of dry litter disposal for the long-term welfare of the communities we live in. The following are the minimum requirements.

Nutrient Management Plan

A good Nutrient Management Plan involves determining the soil requirements, the nutrients in the litter, and applying the proper nutrients at agronomic rates for the crop intended. The Plan should demonstrate methods of litter usage as a fertilizer that maximizes plant uptake and minimizes adverse impacts. Best Management Practices also dictate such things as the development of buffer zones between bodies of water and the application sites and rotation of application fields to prevent buildups of any particular mineral. Each Nutrient Management Plan will, therefore, be site-specific.

Most mandatory regulations require that Nutrient Management Plans be updated when the animal population increases by 10% or more, whenever there has been a significant change in the operations, or no less than every five years. The Nutrient Management Plan should





include manure application volumes, application sites, the crop to be using these nutrients, and timing of the application throughout the year.

These Plans must be prepared by a person certified through the Natural Resources Conservation Service. To develop a valid Nutrient Management Plan, you should contact the Natural Resources Conservation Service, Soil & Water Conservation District, Cooperative Extension Service, or in some cases, a certified private engineer.

Soil Samples

Soil from fields receiving applications of poultry litter should be sampled for a standard soil fertility analysis. The analysis should be utilized to manage the nutrients in the soil, which will prevent a build-up of nutrients that could result in adverse water quality. The soil should be sampled for nitrogen, phosphorus, and potassium content at a minimum. The Producer should then rotate application fields having nutrient concentration to prevent any build-ups of mineral.

A representative soil sample from each litter application site should be taken and analyzed annually. It is recommended that all soils be sampled prior to application. The Cooperative Extension Service should be available to conduct the laboratory analysis and to provide information on the proper time and method to collect the sample.

Producers should retain the lab results for a minimum of three years.

Litter Samples

Litter samples should be utilized to determine the fertilizer potential of the litter and to calculate the application rate. The litter should be sampled for nitrogen, phosphorus, and potassium content at a minimum. In some states, a standard or average nutrient content of the litter has been determined through extensive study and is acceptable. However, these standards or averages are typically very broad, and with the proper documentation of your own representative litter samples, you may be able to land apply more than the standard would allow.

Samples should be taken from each house in such a way that they are representative of the litter throughout the house. These should be collected and analyzed annually. The Cooperative Extension Service should be available to conduct the laboratory analysis and to provide information on the proper method to collect the sample.

Producers should retain the lab results for a minimum of three years.







Litter Cleanout & Application

Effective utilization of manure is possible if you know how much is being spread over a given area. Therefore, litter application equipment should be calibrated prior to usage to ensure the volumes detailed in the Nutrient Management Plan to improve utilization of the nutrients in the manure and reduce the potential for nutrient runoff. Manure should not be applied when the soil is saturated, covered with ice or snow, during precipitation, or when significant precipitation is expected within 24 hours.

Manure should be land applied immediately after its removal from the poultry houses. Any manure stockpiled for later use must be covered in a manner to prevent contact with precipitation. The preferred method of storage would be in a covered dry stack storage structure. The bottom or base of the storage structure should be constructed of either concrete or impermeable clay to prevent ground water contamination.

Buffer zones and vegetated filter strips, as detailed in the Nutrient Management Plan, should be maintained during application. Buffer zones should be no less than 100 feet from all water sources or 50 feet from neighboring occupied dwellings, except where state law dictates otherwise. Most states or counties will have additional guidelines available through the Cooperative Extension Service.

Producers should maintain litter application records detailing the volume, application rate, the acreage covered, and the date of all litter applications. The site or field of manure application should also be recorded, and should only be made on soils detailed in the Nutrient Management Plan. Records should be available for review and be retained for a minimum of three years.

Managing Dry Poultry Litter to Prevent Spread of Disease

While environmentally sound management of poultry litter is important, there are also potential impacts on poultry health, related to movement and spreading of dry poultry litter. Producers realize the importance of disease control and biosecurity related to managing poultry farms, but managing dry poultry litter in a biosecure manner is a tool that can be used to prevent or reduce disease spread.

Even when disease problems are not present in flocks where litter is being removed from houses, litter should be removed from farms only on covered trucks and spread well away from other poultry operations.

Litter from poultry houses where disease problems are present is a potential source of disease infection for other poultry. Used litter contains insects and other poultry disease-causing organisms. Spreading this litter can spread insects and diseases from one location to another. Measures should be taken in all instances to reduce the potential for disease spread, but especially if diseases such as Laryngotracheitis Virus (LT), Avian Influenza (AI), or other infectious transmittable diseases occurred in flocks that were raised on the litter being







removed from a house. Handling dry poultry litter in a responsible manner as related to disease prevention and control becomes even more critical under these circumstances.

Sale of Dry Poultry Litter

Since the generator of the litter is ultimately responsible and liable for the litter if any litter mishandling occurs, Producers selling or giving away the manure from their operations should ensure the landowner receiving the manure has implemented a site-specific Nutrient Management Plan prior to application as well. The Producer should supply the landowner a copy of the most recent litter analysis and maintain records of all sales or transfers of manure, which will include the name and address of the buyer and location of the application site. In addition, an agreement form to transfer or assign the responsibility for the litter's proper utilization may be utilized to help provide a paper trail in the event of future regulatory action.

Training

Producers should attend voluntary training programs on nutrient management, which Tyson will coordinate with the Natural Resources Conservation Service or a similar agency in your area. In most states, the local Cooperative Extension Service has also developed information pamphlets that can help Producers better manage manure application. The information includes best management practices to be utilized for the most economical and practical use of their litter.

Compliance

Because of the critical nature of managing dry litter disposal, Tyson must expect its Producers to comply with "Best Management Practices." If repeat violations of the "Best Management Practices" are documented, the Company will withhold placement of birds until an agreement to comply is obtained.







Dead Bird Disposal

Regulations

Federal, State, and County regulations are also becoming more prevalent in dead bird disposal. Where these regulations are in place, it is our responsibility to follow them completely.

Regardless of regulatory requirements, it is Tyson's expectation that we minimize the environmental impact of dead bird disposal through the utilization of Best Management Practices for the long-term well being of the communities we live in. Approved methods for dead bird disposal vary by state.

Methods

Acceptable methods for disposal of dead birds include:

Composters
Incinerators
Digesters
Freezers (store birds for rendering)
Rendering – Tyson approved
Pits - Not acceptable in many states. Where allowed, must follow state guidelines.
Burial – Available in some states in catastrophic circumstances.

It is unacceptable to leave dead birds piled outside the poultry house where dogs and varmints can carry them off. This practice leads to a concentration of varmints, and the carcass remains spread by dogs or varmints will be objectionable to neighbors. Obviously, birds of prey feeding on dead carcasses also spread unwanted diseases.

Approved methods for disposal of dead birds vary from state to state. Your Technical Advisor can advise you on the approved methods of dead bird disposal for your state.

Compliance

If unacceptable dead bird disposal methods are observed, or if there are complaints from neighbors or regulatory agencies, Tyson will ask the Producer to utilize Best Management Practices. If acceptable practices are not utilized, Tyson will withhold placement of birds until a satisfactory agreement is reached.





Case 4:05-cv-00329-GKF-PJC Document 2120 Filed in USDC ND/OK on 05/29/2009 Page 11 of 75 NUTRIENT MANAGEMENT



NUTRIENT MANAGEMENT PLANS

Land application is the best use of the nutrients in poultry manure. It is relatively inexpensive and is environmentally safe if managed properly. To protect against overapplication, the nutrient concentration of the manure must be known. Application rates can be calibrated to maximize plant nutrient uptake when manure nutrient concentrations, soil fertility, and crop nutrition requirements are known.

A site-specific nutrient management plan can consist of the following:

- -- farm site maps
- -- projected crop yields
- -- soil and manure testing results
- -- buffer zones and filter strips
- -- application rates based on a limiting nutrient
- -- Best Management Plans
- -- method(s) to calibrate application equipment

The Natural Resources Conservation Service, Cooperative Extension Service, and other agricultural research agencies have prepared tables of the average concentrations or amounts of key nutrients found in many animal manures. Tables are included with this handbook for the estimation of the nutrient content of poultry manure. Site-specific testing results, though, are the best source of information to prepare the Nutrient Management Plan.

Once the Nutrient Management Plan has been implemented, it should be reviewed and updated at least every five years to ensure the Plan's effectiveness. The Plan can save the producer money be reducing the amount of commercial fertilizer which may be purchased.

Land Application

One of the most important elements of land application is to apply manure during the plant growing season. The amount of manure applied, the timing of the applications, and the application method used, combine to maximize the benefits of the nutrients.

Poultry manure broadcast upon the surface will lose a significant portion of its volatile nitrogen compounds. Poultry manure applied to soils which are frozen or snow covered has a much higher potential for runoff to surface water sources. Application should not be made proximal to water wells, springs, or sinkholes or on hillsides with slopes greater than 15%.

Incorporation of the manure into the soil increases the amount of available nutrients and reduces the potential for adverse impacts to water quality. Incorporation







results in little disturbance to the soil surface and is appropriate for till and no-till systems.



POULTRY NUTRIENT MANAGEMENT

The majority of producers will apply poultry manure to their pastures or croplands which will be used as fertilizer. To maximize fertilizer usage, growers must develop and implement Nutrient Management Plans before the manure is actually removed from the poultry houses. The Management Plan can protect water quality and will provide for economic returns to the producer.

To protect our natural resources and yet remain profitable, poultry producers must utilize the most up-to-date and accurate site-specific sampling data to develop their Nutrient Management Plans. The Natural Resources Conservation Service is a grower support agency that can develop the Plans for growers. Growers can use one the following models to implement an "unofficial" management plan while the Natural Resources Conservation Service develops the site-specific Plan.

Nutrient management incorporates the nitrogen, phosphorous, and potassium content of the manure with the nutrient requirements of the crops growing in the application field. Proper planning prior to manure application, reduces the wasting of valuable nutrients. Proper nutrient management minimizes adverse impacts which can result from nutrient over-application. Accurate nutrient planning can also reduce the need for commercial fertilizer.

To obtain the most benefit from manure nutrients, while eliminating surface and groundwater contamination, the following should be implemented by the Producers:

- -- apply for an Animal Waste Management Plan
- -- determine the limiting nutrient for application
- -- analyze manure and application site soils annually
- -- match manure application rates to meet plant requirements
- -- calibrate waste application equipment prior to usage
- -- incorporate manure into the soil, if required
- do not apply manure to frozen or saturated soils
- -- apply manure during the active plant growing season
- store manure properly prior to application
- -- establish and maintain filter strips between application sites and water sources.

Training, technical assistance, and financial aid is available to help growers identify and correct problems using poultry manure. The Natural Resources Conservation Service and Cooperative Extension Service are two grower support agencies that can provide assistance.





Page 15 of 75

NUTRIENT MANAGEMENT PLAN DEVELOPMENT

Producers must develop Nutrient Management Plans to prevent excess soil nutrient accumulations which could result in surface and ground water contamination. The Plan must match the crop nutritional requirements with the nutrients available in the manure. The value of the manure depends on its nutrient composition and application practices.

The purpose of the Nutrient Management Plan is to determine the number of acres of pasture or cropland needed for the proper manure application. The Plan will usually have nitrogen or phosphorous as the application limiting nutrient. A limiting nutrient is usually that which will have the greater adverse environmental impacts if application is mismanaged. Plan development requires realistic crop yields in the application areas, their nutrient requirements, and proper timing of application for maximize plant use.

The following information can be used to develop and utilize a Nutrient Management Plan:

- 1. Obtain aerial photographs for each field identified to receive manure
- 2. Obtain a soils survey booklet from the Natural Resources Conservation Service
- 3. Determine fields available for land application
- 4. Realistic Yield Expectations from soil survey or site-specific data base
- Litter nutrient composition from one of the site-specific litter analysis or one of the enclosed tables.

Example One:

Follow the enclosed "Worksheet for Crop Nutrient Requirement" and Tables 1 and 2 to develop a Nutrient Management Plan for your farm. The example uses nitrogen as the limiting nutrient for the calculation manure application rates.

Example Two:

Tables 3 has been added along with another sample worksheet from Example One to determine phosphorus application rates. Note: the resultant application rate is for nutrient uptake only. Phosphorus based application rates are usually calculated on the soil's capacity to assimilate phosphorus rather than on plant phosphorus uptake rates. This is the reason why soil samples are so important. Sampling can alert the producer to excess phosphorous accumulation in application site soils.





Example Three:

A third example entitled "Determine Rate of Broiler Litter Application" is enclosed for reference. This example will help the producer determine phosphorous application rate.

Finally, a worksheet for calculating the volume of manure produced at the farm is included. While worksheets will provide an estimated quantity of litter being generated, poultry operations should establish their own records. The quantity of litter being removed from a poultry house can be determined by litter spreader calibration and calculating the number of loads from the poultry houses.





Worksheet for Calculating Poultry Litter Production

The total amount of litter removed from farm production facilities can be estimated by the following method. The amount is somewhat dependent on the type and amount of litter added initially and may vary by as much as 20% depending on the farm practices.

Broiler House Litter	Example Actual
a. Number of birds per flock:	25000
b. Number of flocks per year:	5.8
c. Pounds market live weight per bird:	4
d. Tons litter per 1000 birds per flock:	1
e. Tons litter per year (a x b x d /1000):	145
Broiler House Manure Cake	Example Actual
a. Number of birds per flock:	25000
b. Number of flocks per year:	5.8
c. Pounds market live weight per bird:	4
d. Tons litter per 1000 birds per flock:	0.34
e. Tons litter per year (a x b x d /1000):	50
Broiler Stockpiled Litter	Example Actual
a. Number of birds per flock:	25000
b. Number of flocks per year:	5.8
c. Pounds market live weight per bird:	4
d. Tons litter per 1000 birds per flock:	1
e. Tons litter per year (a x b x d /1000):	145
Broiler Roaster House Whole Litter	Example Actual
a. Number of birds per flock:	12500
b. Number of flocks per year:	4
c. Pounds market live weight per bird:	8
d. Tons litter per 1000 birds per flock:	2.6
e. Tons litter per year (a x b x d /1000):	130





Broiler Roaster House Whole Litter	Example	Actual
a. Number of birds production facility capacity:	8000	
b. Days per year birds in production facility	365	
c. Pounds market live weight per bird:	6	
d. Pounds litter per 1000 birds per day:	131	waxaanaana
e. Tons litter per 1000 bird capacity per year: (b x d /2000)	24	
f. Tons litter per year (a x e /1000)	192	
Turkey Brooder House Whole Litter	Example	Actual
a. Number of birds per flock:	20000	
b. Number of flocks per year:	7	
c. Average pounds market live weight per bird:	2.5	
d. Tons litter per 1000 birds per flock:	0.75	
e. Tons litter per year (a x b x d /1000):	105	
Turkey Grower Hen House Whole Litter	Example	Actual
a. Number of birds per flock:	12000	1 <u>-1,</u>
b. Number of flocks per year:	3.4	
c. Pounds market live weight per bird:	16	
d. Tons litter per 1000 birds per flock:	5	
e. Tons litter per year (a x b x d /1000):	204	
Turkey Grower Hen House Manure Cake	Example	Actual
a. Number of birds per flock:	12000	
b. Number of flocks per year:	3.4	
c. Pounds market live weight per bird:	16	
d. Tons litter per 1000 birds per flock:	1.7	
e. Tons litter per year (a x b x d /1000):	70	L.





•	

Turkey Grower Tom House Whole Litter	Example	Actual
a. Number of birds per flock:	8000	
b. Number of flocks per year:	3.4	
c. Pounds market live weight per bird:	25	-
d. Tons litter per 1000 birds per flock:	7.5	
e. Tons litter per year (a x b x d /1000):	204	-
Turkey Grower Tom House Manure Cake	Example	Actual
a. Number of birds per flock:	8000	
b. Number of flocks per year:	3.4	
c. Pounds market live weight per bird:	25	
d. Tons litter per 1000 birds per flock:	2.5	
e. Tons litter per year (a x b x d /1000):	70	
Turkey Stockpiled Litter	Example	Actual
a. Number of birds per flock:	8000	
b. Number of flocks per year:	3.4	
c. Pounds market live weight per bird:	25	
d. Tons litter per 1000 birds per flock:	6.6	
e. Tons litter per year (a x b x d /1000):	180	-
Turkey Breeder House Whole Litter	Example	Actual
a. Number of birds production facility capacity:	6000	
b. Days per year birds in production facility:	365	
c. Pounds live weight per bird:	20	Note the Party of
d. Pounds litter per 1000 birds per day:	204	
e. Tons litter per 1000 bird capacity per year: (b x d /2000)	37	
f. Tons litter per year (a x e /1000):	222	





EXAMPLE ONE: NITROGEN

one worksheet per crop.	Example	Your Farm
		,
Crop to be grown	Corn	
Crop realistic yield exceptions from Tables 3 or 4 or on farm records.	110 bu/acre	nak 24 tani na manana mana
Nitrogen fertilization guidelines per unit of yield based on soil type (see Table 1)	1.0 lb N/bu	
Crop Nitrogen requirement (2 X 3)	110 lb N/bu	
Starter fertilizer nitrogen (if needed)	25 lb N/acre	
Commercial fertilizer (if needed)	0 lb N/acre	
Crop Nitrogen need from poultry dry litter 4 minus 5	85 lb N/acre	
Poultry dry litter Plant-available Nitrogen a). Total nitrogen composition of litter from on farm sampling or enclosed Tables 2 or 4	38 lb N/acre	
b). Nitrogen availability coefficient (see Table 2) based on application method	0.5	
c). Plant-avaliable nitrogen (a x b) or from Waste analysis	19 lb N/ton	
Poultry dry litter application rate (7 divided by 8.c.)	4.5 ton/acre	
. Acres of crop to be grown	95 acres	
. Total litter required to meet the agronomic N requirement for this field (9 x 10)	427.5 tons	







Table 1. Crop Nitrogen Factors for Calculating Annual Realistic N Requirements per acre.

Crop	Suggested Nitroge	n Application Rates
	Loamy, Clayey Soils	Sandy, Leachable Soils
Crops Harvested for		
Grain or Fiber		
Corn (grain)	1.0 lb N per bu	1.25 lb N/bu
Sorghum (grain)	2.0 lb N per cwt	2.5 lb N/bu
Soybean (grain)		
(Manure Utilization Only)	3.8 lb N per bu	4.0 lb N per bu
Cotton	0.06 lb N per lb lint	0.12 N/lb lint
Wheat (grain)	1.7 lb N per bu	2.4 lb N/bu
Barley (grain)	1.4 lb N per bu	1.6 lb N /bu
Oats (grain)	1.0 lb N per bu	1.3 lb N/bu
Rye (grain)	1.7 lb N per bu	2.4 lb N/bu
Triticale (grain)	1.4 lb N per bu	1.6 lb N/bu
Silage, Hay, and		
Pasture Crops		
Corn (silage)*	10.0 lb N per ton	12.0 lb N per ton
Sorghum-sudangrass*	45.0 lb N per ton	55.0 lb N per dry ton
Bermudagrass*	40.0 lb N per ton	50.0 lb N per ton
Tall fescue*	40.0 lb N per ton	50.0 lb N per dry ton
Orchardgrass*	40.0 lb N per ton	50.0 lb N per dry ton
Timothy*	40.0 lb N per ton	50.0 lb N per dry ton
Annual ryegrass*	40.0 lb N per ton	50.0 lb N per dry ton
Small grain*	50.0 lb N per ton	60.0 lb N per dry ton
Millet*	45.0 lb N per ton	55.0 lb N per dry ton
	71 <u>. – 1</u> 10. <u>– 110. – </u>	
Forest Species		
Pines**	40.0 lb N per year	60.0 lb N per year
Hardwood trees* *	70.0 lb N per year	100.0 lb N per year

^{*} Reduce N rate by 25 percent when grazing.





^{**} On trees less than 5 feet tall, nitrogen will stimulate undergrowth competition.



Table 2. Average nutrient content of litter and N availability factors by production system and handling method.

		N Availability Factor	lity Factor		
	Total N	Broadcast	Incorporated	Total P2O5	Total K20
	lb/ton	0.5xlb.T.N/ton	0.6xlb.T.N./ton	lb/ton	
Poultry House					
Litters					
Chicken					
broiler	72.3	36.15	43,38	78.5	45.8
roasfer	73.3	36.65	43.98	74.8	44.9
breeder	31.2	15.60	18.72	54.0	31.0
Turkey					
poult	45.3	22.65	27.18	52.0	32.1
grower hen	57.2	28.60	34.32	72.5	40.4
grower tom	57.2	28.60	34.32	72.5	40.4
Poultry					
Stockpiled				19.00p.00.000	
Litters					
Broiler	35.6	17.80	21.36	79.6	34.5
Turkey	35.7	17.85	21.42	72.3	33.0



EXAMPLE TWO: PHOSPHORUS

orksheet: Crop Nutrient Requirement Wor	ksheet.	
se one worksheet per crop.	Example	Your Farm
1. Crop to be grown	Corn	
Crop realistic yield exceptions from Tables 3 or 4 or on farm records.	80 bu/acre	
3. Crop Phosphorus requirement from Tables 3 or 4	42 lb P/acre	
4. Commercial fertilizer (if needed)	0 lb N/acre	
5. Crop Phosphorus need from poultry dry litter (3 minus 4).	42 lb P/acre	William Communication of the C
6. Total phosphorus composition of litter from farm sampling, or see Table 2 or 4.	82 lb P/ton	The second secon
7. Poultry dry litter application rate (6 divided by 5)	0.5 ton/acre	
8. Acres of crop to be grown	95 acres	
Total litter required to meet the agronomic phosphorus requirement for this field (8 x 9)	50 tons	





Table 3. Crop Nutrient Utilization.Source: Potash Phosphate Institute of America.

l	S.	1	
_	_		

Crop	Yield	N	P2O5	K20
			lb/acre	
Corn	. 80 bu	121	42	77
	100 bu	160	60	120
	150 bu	1 8 5	80	215
	180 bu	240	100	240
,	100 00			
Corn Silage	16 tons	130	45	102
oom onago	32 tons	200 ·	80	245
	02 to113	200		
Soybeans	30 bu	123	32	52
ooybeans	40 bu	180	45	80
	50 bu	257	48	120
	60 bu	336	65	145
	90 DU	330	00	7.10
Grain Sorghum	4 tons	250	90	200
Gram Sorgitum	4 (0113	200	•	200
 Wheat	40 bu	70	30	50
e atteat	60 bu	125	50	110
1	80 bu	186	54	162
	60 Du	100	04	102
Oats	80 bu	75	35	95
Cals	100 bu	150	55	150
	100 00		4.5	
Barley	65 bu	74	32	63
Durisy	100 bu	150	55	150
	100 00	,,,,	-	
Alfalfa	4 tons	180	40	180
rinana	8 tons	450	80	480
	0 (0)13	.00		
Orchardgrass	6 tons	300	100	375
Brome grass	5 tons	166	66	254
Tall fescue	3.5 tons	135	65	185
Blugrass	3 tons	200	55	180
Costal Bermuda	O LONG	200	V 5	100
grass	4 tons	225	. 40	160
grass	10 tons	535	145	410
Claver	4.5 tons	185	60	175
Clover grass		300	90	360
	6 tons	300	an	300
Guerr honte	30 tons	275	85	550
Sugar beets	30 10118	210		000
Rice	2.25 tons	110	45	110
11100	3.5 tons	112	60	168
Timothy	4 tons	150	55	250
Timothy		299	108	430
Panola grass	12 tons	299	100	450
Sorghum-Sudan				
grass	8 tons	319	122	467
G1893	0 10110	<u> </u>		



from MWPS-18, Midwest Plan Service, Iowa State University, Ames, IA.



EXAMPLE THREE

DETERMINE RATE OF BROILER LITTER APPLICATION

1. Determine nutrient recommendation (lbs/ac) for crop to be grown from either Table 3 or 5:


Сгор			
Nutrient requirement (Lbs/Ac)	N	P2O5	K20
2. Soil Test	-		hadden bereit and the state of
3. Broiler Litter Analysis: (Use Table 4 if analysis is not available.)		***************************************	
4. Subtract the nitrogen in the		en the crop requires	3 .
N crop requires N in soil (Soil test PPM X 2) Total N required	N 	-	
 Determine rate of broiler litte N crop requires = Rate/litter, T N in litter 		;) 	
6. Nutrient balance:	N	P2O5	K2 0
Pounds nutrients required by crop (#4)			**************************************
Pounds nutrients applied intons litter (No. 5 x No. 3)			<u> </u>
(ito, o it ito. o)	***************************************		



7. Determine if the amount of phosphorus applied is greater than twice the amount of phosphorus needed by the crop or grass. If this is true, the rate of broiler litter should be reduced one ton per acre, or until the amount of phosphorus applied is no greater than twice the amount of phosphorus required by the crop should be applied.

		N	P2O5	KZ
Nutrient Bala	nce:			
Pounds nutri	ents			
required by c	rop:		***************************************	
Pound nutrie	nt supplied			
by			****	
Balance:		P. C.		April 1990
The amount			o meet crop requiren	nents after
	TODS OF BEOREE DE	ter is applied.		
	_ tons of broiler lit	ter is applied.		





		Nutrient Reco	mmendation	
Crop	Yield Goal	N	P2O5	K20
Corn	75-99 bu/acre	75-100	60	20
	100-149 bu/acre	110-165	80	130
	150-200 bu/acre	180-240	80	140
Cotton	1.0 bale/acre	40	40	30
	1.5 bales/acre	60	60	50
	2.0 bales/acre	80	80	80
	2.5 bales/acre	100	80	80
Grain Sorghum	1500-2000 lbs/acre	30-40	20	20
	2000-4000 lbs/acre	40-80	40	80
	4000-6000 lbs/acre	80-120	60	100
	6000-8000 lbs/acre	120-160	80	120
Wheat	20-30 bu/acre	40-60*	20	20
	30-40 bu/acre	60-80	40	30
	40-60 bu/acre	80-120	40	40
	60-80 bu/acre	120-160	60	60
	80-100 bu/acre	160-200	60	60
Coastal Bermuds	grazing only	100-160	50	90
	1 cutting + grazing only	160-220	50	150
	3 cuttings	300-350	100	300
	4-6 cuttings	400-600	130	40 0
Alfalfa	non-irrigated, annually	20	60	120
	irrigated> 6 T/acre	20	100	120
	irrigated > 8-12 T/acre	20	140	200
Clover	annually	20	80	120
	sod seeded	20	80	120
	with ryegrass/small grain	40	80	120
Wheat	light grazing**	160	60	60
	moderate grazing	200	80	120
	heavy grazing	240	80	120
Sorghum/Sudan	1 cutting/light grazing	80	40	40
	2 cuttings/medium grazing	160	60	60
	3 cuttings/heavy grazing	200	80	80

Source: Texas Agricultural Extension Service Soil Testing Laboratories, College Station and Lubbock.



Case 4:05-cv-00329-GKF-PJC Document 2120 Filed in USDC ND/OK on 05/29/2009 Page 28 of 75

SOIL AND MANURE SAMPLINGS



APPLICATION SITE SOIL SAMPLING

All manure application fields which receive poultry manure should be sampled annually. The Nutrient Management Plan will include a site map of all application fields which should be sampled. Soil samples should be collected and submitted to the testing laboratory prior to the end of April or at least one month prior to the beginning of the active growing season. The sampling results will detail nutrient and mineral concentrations of the soil. The sampling laboratory will usually include fertilizer recommendations to further maximize crop nutrient needs along with the sampling results.

Sampling Laboratory


Soil analysis is usually provided by the State Cooperative Extension Service. Private soils testing laboratories can also be utilized for analysis. The local county Extension Office or private lab should be contacted for details on sampling methods and recommended analysis. Generally, the Extension Office or laboratory will provide sampling instructions, information sheets, and sampling boxes. Depending on the state, a nominal fee is sometimes charged for analysis done by the state. Private laboratory fees can vary. Each sample should be analyzed for, at a minimum, nitrogen, phosphate, and potassium. The Cooperative Extension Service or sampling laboratory will usually provide information or consultation to explain the results of the analysis.

Collecting Soil Samples

To collect samples, a small volume of the soil should be taken from approximately 20 random locations in each application field. The enclosed diagram illustrates several different sized fields and suggestions for sampling patterns. Soil should be collected from a depth of four to six inches below the surface at sampling location. The small samples should then be thoroughly mixed in a plastic bucket. A portion of this larger sample should then be air dried and poured into the provided sample box and submitted to the Extension Service or soil testing laboratory.







Sampling patterns for different field sizes



POULTRY LITTER SAMPLING

Prior to application, a representative poultry manure sample should be taken and analyzed. The nutrient concentration of the manure will allow the producer to accurately calculate application rates according to the Nutrient Management Plan.

Sampling Laboratory


The local Cooperative Extension Service is usually available to conduct analysis of the manure. Also, a private laboratory can be used for analysis. Please contact the sampling laboratory prior to actually collecting the sample. The sampling laboratory will usually provide a sample identification form. At a minimum, the manure should be sampled for nitrogen, phosphorous, and potassium. It is recommended that calcium, magnesium, sulfur, iron, manganese, zinc, and copper be included in the analysis. The Cooperative Extension Service or sampling laboratory should also provide information or consultation to explain the results of the analysis.

Sample Collection



The sample should be composed of approximately 20 smaller samples taken from random locations within each house. The enclosed diagram illustrates potential sampling locations. The samples should be representative of the clean-out material. Cake litter samples should also be taken and analyzed prior to its removal and application. Mix the smaller samples in a plastic bucket. Pour a portion of the thoroughly mixed sample into a quart size plastic bag. Close the bag and write farm and house identification information and submit to the Extension Service or private laboratory.





NOTE: FROM THE NORTH GAROLINA "DRY POLLTRY LITTER HANDSOOK"

Case 4:05-cv-00329-GKF-PJC Document 2120 Filed in USDC ND/OK on 05/29/2009 Page 33 of 75

RECORD KEEPING FORMS





RECORD KEEPING FORMS

The sheets in this section will help the producer keep field application records, field application rates, and also help with determining the amount of manure taken from his individual houses.

Record keeping provides a convenient reference that allows the producer to maximize manure fertilizer usage. The records also ensure that the volume of manure applied is following the Nutrient Management Plan. The following sheets will allow the grower to "follow" the manure from when it leaves the individual house to the application site. This is especially useful when manure is sold or given away.





Other*** (tons) from farm Removed (tons) Use of Litter . . Spread on Farm (tons) Temporary Storage (tons) 3rd Party Applicator 3rd Party's Address 3rd Party's Phone # Dry Litter Cleanout Records Total Removal (tons) (4) x (2) Spreader Capacity** (tons) Spreader Volume* (cubic feet) # of loads removed Owner's Address Owner's Phone # Farm Owner mm/dd/yy Date

. Can be found in operator's manual for the spreader. Contact a local dealer if you do not have your owner's manual.

FOTALS

:

** See attached sheet for formula.

*** Indicate use (i.e., catlle feed, composted soil ammendment, commercially distributed organic fertilizer, etc.)

For Recording Dry Litter Application Events on Different Fields Dry Litter Application Field Record


Spreader Operator Farm Owner

Spreader Capacity*** (tons)	
# of Loads Volume of Loads** per Field (cubic feet)	
# of Loads	
Application Method*	
Field Size	
Crop Type	
Date (mm/dd/vy)	
Field #	
Tract #	

· SI= Soil incorporated (disked); BR= broadcast (surface applied)

"Can be found in operator's manual for the spreader. Contact a local dealer if you do not have your owner's manual.

***See attached calibration sheet for the formula.



^{***}Enter the value received by subtracting column (7) from (B). Continue subtracting column (7) from column (B) following each application event. ** See your waste management plan for sampling frequency. At a minimum, waste analysis is required within 60 days of land application events. · Can be found in operator's manuel for the spreader. Contect a local dealer if you do not have your owner's manual,



ANIMAL WASTES APPLICATION RECORDS

PRODUCER'S NAME:_	
WASTE APPLICATION	YEAR:

FIELD NO.	LAND OWNER	DATE	ACRES COVERED	APPROX. VOLUME APPLIED
	<u> </u>			





CALIBRATION OF APPLICATION EQUIPMENT

Spreader Calibration

Proper calibration of manure spreading equipment controls the amount or volume applied. Calibration ensures the spread of nutrients according to the Nutrient Management Plan. Poultry growers can calibrate their equipment by using site-specific soils, manure, and plant data which maximizes economic returns of land application.

Technical assistance is available from the Natural Resources Conservation Service, Cooperative Extension Service, and other Federal, State and local producer support agencies. Information sheets from these agencies are usually available which will help growers determine spreader capacity and application rates. Below are examples to help producers calibrate spreader equipment.

Calibration

To begin calibration, locate an open area which is fairly smooth. Spread a large plastic sheet or tarp over the ground. The following table "Calibrating of Manure Spreaders" uses tarp sizes of 8' x 8', 10' x 10', and 10' x 12' for calibrations. Fill the spreader and drive it over the tarp as it is applying manure. To collect a representative sample, the manure spreader should be driven at least three times over the tarp. Next, use a scale and weigh a plastic bucket. Record the weight. Pour the manure collected into the bucket and weigh. Subtract the weight of the bucket from this weight. This will then be the weight of the manure. The producer should repeat the above calibration procedure three times and then average the weight of the material applied.

Calculation of Application Rate

The following table can be used to determine application rate if the weight of the sample and size of the tarp are known. The application rate can also be determined by using the formula below:

Application Rate (tons/ac) = weight of manure x 21.78 tarp length x tarp width

The application rate of the manure can also be adjusted by changing the speed at which the manure is applied. The following formulas can be used for calibration. To correctly utilize the formulas, the producer must know: the spreader load capacity in tons, application time in minutes, and the average width of the manure spread from the applications in feet.

Application Rate (tons/ac) = $\frac{\text{spreader load x 495}}{\text{time x width x speed}}$





Travel Speed (mph) = $\frac{\text{spreader load x 495}}{\text{time x width x application rate}}$

Spreader Pattern

The uniformity of the manure spread and determination of spreader overlap can easily be determined. To do so, place several small sized approximately 12" x 12" or 15" diameter pans, 2' to 4' across the spreader path. Next, drive the spreading equipment through the middle of the pan arrangement. Weigh the manure in each of the pans or visually compare the amounts in each pan. The width of the manure spread can be determined by comparing the volume or the lack of material in each pan. Once the width is determined, application can overlap the edges to ensure uniform coverage.








CALIBRATING OF MANURE SPREADERS

SIZE OF PLASTIC SHEET OR TARP					
Pounds of Manure	8' X 8'	10' X 10'	10' X 12'		
Applied to Sheet	Tons of Manure Applied Per Acre				
1	0.34	0.22	0.18		
2	0.68	0.44	0.36		
3	1.02	0.65	0.54		
4	1.36	0.87	0.73		
5	1.70	1.09	0.91		
6	2.04	1.31	1.09		
7	2.38	1.52	1.27		
8	2.72	1.74	1.45		
9	3.06	1.96	1.63		
10	3.40	2.18	1.82		
11	3.74	2.40	2.00		
12	4.08	2.61	2.18		
13	4.42	2.83	2.36		
14	4.76	3.05	2.54		
15	5.10	3.27	2.72		
16	5.45	3.48	2.90		
17	5.79	3.70	3.09		
18	6.13	3.92	3.27		
19	6.47	4.14	3.45		
20	6,18	4.36	3.63		
21	7.15	4.57	3.81		
22	7.49	4.79	3.99		









BEST MANAGEMENT PRACTICES

Best Management Practices (BMP's) have been utilized on poultry farms for several years. The BMP's address manure handling practices, filter strips, soil and litter testing, and record keeping. The following is a discussion of different management practices which may be applicable to poultry producers.

Composting

Composting can be used to biologically stabilize manure into a beneficial soil amendment. Producers can mix crop residuals, by-products, and other "waste" materials with manure for composting. Composting reduces odor and flies. Heat generated during composting destroys harmful organisms which may be within the manure. The volume of the final compost product is less than the volume of the initial manure, saving the producer hauling and spreading expense.

Field Buffer Zones

Field borders are vegetated strips which are established at the edge of application sites. The buffers reduce erosion and the runoff of manure from the application site. The reduction of nutrient runoff enhances overall water quality.

Riparian Buffer Zones

Grass or forest filter strips can protect water quality proximal to poultry farins and application sites. The vegetated strips act to filter excess nutrients from runoff. The plants in the filter strip then utilize the nutrients for their nutritional needs. The strips do remove acreage from production, but the reduction of adverse impacts to water quality demonstrates good environmental stewardships. Contact the local Natural Resources Conservation Service, Cooperative Extension Service, or other grower support agency for additional details.

Acceptable Best Management Practices

The following is a general listing of Best Management Practices from the Texas Agricultural Extension Service which can reduce adverse water quality impacts from poultry operations:

- 1. All litter stockpiled or retained on site shall:
 - a. Be stored under cover to be protected from rainfall
 - b. Be placed on impermeable clay or concrete to prevent infiltration
 - c. Be isolated from all run-off waters by dikes, terraces, berms, ditches or other structures





- Maintain grass filter or barrier strips between manure application fields and streams or lakes, and other water sources.
- 3. Do not spread broiler litter within 150 feet of any water well, or 100 feet from a highway and neighboring property line.
- 4. Do not apply litter to land with slopes greater than 15 percent.
- 5. Soil test 4-6 inches deep each year and maintain results.
- 6. Sample litter before application
- 7. Cover trucks hauling litter more than one mile.
- 8. Do not apply litter to frozen or saturated soils or during rain or snowfall.
- 9. Do not apply litter to highly erodible lands.
- 10. Inform neighbors when you will be applying litter.
- 11. Apply manure early in the week. Avoid spreading on Fridays or during the weekend.
- 12. Maintain records:
 - a. Where litter was applied
 - b. How much litter applied per acre
 - c. When litter applied and to what crop
 - d. Anyone you sold or gave litter to
 - e. Location where litter will be applied by producer you sold or gave litter to.

Odor Control

The following table is of Best Management Practices to control odors which may result from poultry operations.





Poultry Farm Waste Management Odor Control Checklist:

<u>Source</u> Farmstead	<u>Cause</u> Poultry Production	BMPs to Minimize Odor Vegetative or wooded buffers Recommended best management practices Good judgment and common sense
Ventilation exhaust fans	Volatile gases Dust	Fan maintenance Efficient air movement
Indoor surfaces	Dust	Vacuum and washdown between flocks
Feeders	Feed spillage	Design, operate and maintain feed system to minimize accumulation of decaying wastage. Clean up spillage on a routine basis
Feed storage	Decomposition of accumulated feed residues	Reduce moisture accumulation within and around immediate perimeter of feed storage by ensuring drainage is away from site and/or providing adequate containment Clean up spillage on a routine basis
Litter storage and handling areas	Decomposition of accumulated manure	Remove spillage on a regular basis Provide for adequate drainage around manure stockpiles Inspect for and remove or break up accumulated wastes in filter strips around stockpiles and manure handling area as needed.
Dead birds	Carcass decomposition	Proper disposition of carcasses
Incinerators	Incomplete combustion	Secondary stack burners



BMPs to Minimize Odor

Carcass decomposition

Cause

Source Dead bird disposal pits

Grade and landscape such that water drains away from facilities

Standing water around facilities Improper drainage
Microbial decomposition of organic matter

Complete covering of carcasses in burial pits Proper location/construction of burial pits Disposal pit covers tight fitting

Farm access road maintenance

Poorly maintained access roads

Mud tracked onto public roads

from farm access

(From the North Carolina "Dry Poultry Litter" Handbook)



Transportation of Poultry Manure

In the U.S., the majority of poultry producers are concentrated within a 40 mile radius of the complex feed mill, hatchery, and processing plant. Producers must monitor their application soils for nutrient build-up. If excess nutrient concentrations are determined by soil sampling, the producer must either increase the number of application sites or transport the manure to other areas. In locations where suitable spreading lands are not available, other uses and markets must be determined and implemented for the manure.

Should more than one poultry production company operate in the same geographic area, then the competition for suitable application acres can be rather intense. Producers could form cooperative agreements with one another to establish markets and alternative uses for their manure. The final users of the product could even be several hundred miles from the poultry farms which generated the manure. Transporting manure out of areas with high nutrient concentrations can greatly reduce the potential for water quality impacts.



The manure from disease free farms should only be that which is transported to outside locations. All trucks used for manure transport should be properly cleaned and disinfected to prevent adverse health impacts. The manure should also be covered to prevent it from being blown out of the truck during transport. If the manure is to be stored on-site while awaiting transport, it must be covered to prevent contact with precipitation. The manure should also be stored in plies to promote the compost process which can control the spread of harmful bacteria.



MANURE STOCKPILES AND STORAGE STRUCTURES

When poultry manure is removed from houses, it is not always possible to land apply immediately. Delays to application can include the time of the year, plant growing season, and inclement weather.

Permanent Structures

A roofed structure is the preferred method to store manure prior to application. Such structures must have their foundations constructed from concrete or of an impermeable clay. This eliminates runoff and leaching to ground and surface waters. The roof eliminates the introduction of excessive moisture which can lead to a reduction of nutrients.

Temporary Structures

Producers can use windrow or bunker type arrangements for temporary manure storage facilities. The temporary structures must also be covered and foundation constructed of impermeable clay or concrete to protect the manure from rainfall and runoff.

Temporary structures should be sited in areas which are well-drained to avoid the collection of rain water. The site should be at least 100' from surface waters and drinking water sources. Filter strips should also be maintained around temporary structures to prevent nutrient runoff.

Preventing Fires in Storage Structure

The following guidelines will help reduce the potential for fires:

- keep the litter dry and away from the end of the barn
- do not mix the wet cake with dry litter
- limit stack height to less than 5 feet
- internal temperature of the stack should not exceed 180 °F.





Case 4:05-cv-00329-GKF-PJC Document 2120 Filed in USDC ND/OK on 05/29/2009 Page 51 of 75

MORTALITY



MORTALITY MANAGEMENT

Producers are beginning to implement environmentally sound measures for disposal of their dead poultry. If not properly disposed, dead birds can cause odor, water quality, disease, insect, rodent, and small animal problems. Proper disposal methods can include rendering, composting, incineration, and disposal pits. Information of disposal methods can be obtained at the local office of the Natural Resources Conservation Service or Cooperative Extension Service.

Rendering Conversion

Rendering is an excellent dead bird disposal method. Almost 100% of the bird can be converted into alternative uses. Rendering removes mortalities from the farm which also eliminates the potential for adverse environmental impacts. Tyson is providing its growers with freezers to chill the dead birds prior to transport to the rendering facility.

Composting



Composting dead birds is another practical and economic method of dead disposal. The final product can be used as a soil amendment or conditioner. Besides the birds, waste materials such as litter, straw, and by-products can be used for compost.

Incineration

One of the major advantages of incineration is the virtual elimination of adverse water impacts. Incinerators, though, need to be carefully sited to reduce possible odor complaints. Poultry growers must address air quality issues such as odor and dust which can be generated by incineration. Producers considering incineration should contact state and local environmental agencies for regulatory measures which must be adopted for design and use.

Disposal Pits

Burial of dead birds in open pits and trenches is not an acceptable disposal practice. States that do allow burial, usually require the installation of fabricated units or boxes. Disposal pits require little labor, but must be properly maintained to prevent nuisance complaints.

Disposal pits can be constructed from several sources which include concrete, concrete blocks, or treated timbers. A septic tank could even be used for the disposal pit. The structure usually has one or more openings to the surface to allow the dead birds to be placed inside. The openings should always be covered to prevent the escape of unpleasant odors.





The disposal pit should be located at least 200' from dwellings, 300' from water sources, and 25' from the poultry houses. The pit should be located in soils with good drainage to prevent ponding of rain. The bottom of the pit should be at least five above the elevation of the highest known water table.





Case 4:05-cv-00329-GKF-PJC Document 2120 Filed in USDC ND/OK on 05/29/2009 Page 54 of 75

GROWER RESOURCE AGENCIES



GROWER RESOURCE AGENCIES

Poultry Water Quality Consortium

- Encourages the use of poultry by-products as resources rather than allowing the material to contribute to water pollution.
- Promotes cooperation and information exchanges between government and industry on water quality and by-product issues.
- Composed of representatives from: Southeastern Poultry and Egg Association, Natural Resources Conservation Service, Tennessee Valley Authority, and the U.S. Environmental Protection Agency.

The Consortium can be contacted at: HB-2C; 1101 Market Street; Chattanooga, TN 37402-2801; (615) 751-7292 or (615) 751-7479 (FAX)

Southeastern Poultry and Egg Association



- Dedicated to the growth of the entire poultry industry.
- Extensive programs keep members informed of regulatory measures and technical advances made to the industry.
- Grants are available to help researchers discover and develop better production methods.

The Association can be contacted at: 1530 Cooledge Road; Tucker, GA 30084; (404) 493-9401 or (404) 493-9257 (FAX)

Natural Resources Conservation Service

- Administers national soil and water conservation programs.
- Provides USDA leadership to assist landowners implementing resource conservation measures.
- Provides technical support to landowners considering manure storage facilities, mortality disposal practices, and nutrient nutrition plans based on soils and crops.



The NRCS usually maintains a local office in most counties in the United States. Consult the local telephone directory under U.S. Department of Agriculture for the number and address.



Cooperative Extension Service

- Provides research based information and technology to producers.
- Partnership with USDA and each states land grant university.

The CES usually maintains an office in each county of all U.S. states. Consult the local telephone directory under local government for the number and address.





DIRECTORY OF STATE AND NATIONAL POULTRY ASSOCIATIONS

Alabama Poultry and Egg Association One Court Square, Suite 315 Montgomery, AL 36104 TEL: (205) 265-2732 FAX: (205) 265-0008

American Poultry Association 26363 South Tucker Road Estacad, OR 97023 TEL: (503) 630-6759

Arkansas Poultry Federation P.O. Box 1446 Little Rock, AR 72203-1446 TEL: (501) 375-8131

FAX: (501) 375-5519

Delaware Poultry Improvement Association RD 2, Box 48 Georgetown, DE 19947 TEL: (302) 856-7303

Delmarva Poultry Industry, Inc. RD 6, Box 48 Georgetown, DE 19947 TEL: (302) 856-9037 FAX: (302) 856-1845

Florida Poultry Federation 4508 Oak Fair Blvd. Suite 290 Tampa, FL 33610 TEL: (813) 628-4551 FAX: (813) 620-4008

Georgia Poultry Federation P.O. Box 763 Gainesville, GA 30503-0763 TEL: (404) 532-0473 FAX: (404) 532-7543



Page 58 of 75



Georgia Poultry Improvement Association P. O. Box 20

Oakwood, GA 30566 TEL: (404) 535-5996

FAX: (404) 535-1948

Georgia Turkey Association P.O. Box 127 Watkinsville, GA 30677-0127 TEL: (706) 769-5668

Indiana State Poultry Association Purdue University 1026 Poultry Science Building W. Lafayette, IN 47907-1026

TEL: (317) 494-8517 FAX: (317) 494-6349

Kentucky Poultry Federation P.O. Box 21829 Lexington, KY 40522-1829 TEL: (606) 257-2694

Mississippi Poultry Association P.O. Box 13309 Jackson, MS 39236-3309 TEL: (601) 355-0248

FAX: (601) 353-3840

225 E. Capitol Ave. Jefferson City, MO 65101 TEL: (573) 761-5610 FAX: (573) 761-5619

Missouri Poultry Federation

National Broiler Council The Madison Building, Suite 614 1155 15th Street, N.W. Washington, DC 20005 TEL: (202) 296-2622 FAX: (202) 293-4005







National Turkey Federation 11319 Sunset Hills Road Reston, VA 22090 TEL: (703) 435-7206 FAX: (703) 481-0837

North Carolina Poultry Federation 4020 Barrett Drive, Suite 102 Raleigh, NC 27609

TEL: (919) 783-8218 FAX: (919) 783-8220

Oklahoma State Poultry Federation P.O. Box 357 Hartshorne, OK 74547 TEL: (918) 297-7219

Pennsylvania Poultry Federation 500 North Progress Avenue Harrisonburg, PA 17109 TEL: (717) 652-7530 FAX: (717) 652-0230

FAX: (717) 652-0230

South Carolina Poultry Federation 1201 Main Street, Suite 1220 AT&T Building Columbia, SC 29201 TEL: (803) 748-1283

TEL: (803) 748-1283 FAX: (803) 748-1294

Tennessee Egg and Poultry Association

P.O. Box 11082

Knoxville, TN 37939-1082 TEL: (615) 974-7351 FAX: (615) 974-7448

Tennessee Poultry Improvement Board P.O. Box 40627 Nashville, TN 37204

TEL: (615) 360-0120 FAX: (615) 781-5309







Texas Poultry Federation 8130 Burnet Road P.O. Box 9589 Austin, TX 78766-9589 TEL: (512) 451-6816

FAX: (512) 454-4221





POULTRY ASSOCIATIONS



DIRECTORY OF STATE AND NATIONAL POULTRY ASSOCIATIONS

Alabama Poultry and Egg Association One Court Square, Suite 315 Montgomery, AL 36104 TEL: (205) 265-2732 FAX: (205) 265-0008

American Poultry Association 26363 South Tucker Road Estacad, OR 97023 TEL: (503) 630-6759

Arkansas Poultry Federation P.O. Box 1446 Little Rock, AR 72203-1446 TEL: (501) 375-8131 FAX: (501) 375-5519

Delaware Poultry Improvement Association RD 2, Box 48 Georgetown, DE 19947 TEL: (302) 856-7303

Delmarva Poultry Industry, Inc. RD 6, Box 48 Georgetown, DE 19947 TEL: (302) 856-9037

FAX: (302) 856-1845

Florida Poultry Federation 4508 Oak Fair Blvd. Suite 290 Tampa, FL 33610 TEL: (813) 628-4551 FAX: (813) 620-4008

Georgia Poultry Federation P.O. Box 763 Gainesville, GA 30503-0763 TEL: (404) 532-0473 FAX: (404) 532-7543





Georgia Poultry Improvement Association P. O. Box 20 Oakwood, GA 30566 TEL: (404) 535-5996 FAX: (404) 535-1948

Georgia Turkey Association P.O. Box 127 Watkinsville, GA 30677-0127 TEL: (706) 769-5668

Indiana State Poultry Association Purdue University 1026 Poultry Science Building W. Lafayette, IN 47907-1026 TEL: (317) 494-8517 FAX: (317) 494-6349

Kentucky Poultry Federation P.O. Box 21829 Lexington, KY 40522-1829 TEL: (606) 257-2694

Mississippi Poultry Association P.O. Box 13309 Jackson, MS 39236-3309 TEL: (601) 355-0248 FAX: (601) 353-3840

Missouri Poultry Federation 225 E. Capitol Ave. Jefferson City, MO 65101 TEL: (573) 761-5610 FAX: (573) 761-5619

National Broiler Council The Madison Building, Suite 614 1155 15th Street, N.W. Washington, DC 20005 TEL: (202) 296-2622 FAX: (202) 293-4005





National Turkey Federation 11319 Sunset Hills Road Reston, VA 22090 TEL: (703) 435-7206 FAX: (703) 481-0837

North Carolina Poultry Federation 4020 Barrett Drive, Suite 102 Raleigh, NC 27609 TEL: (919) 783-8218 FAX: (919) 783-8220

Oklahoma State Poultry Federation P.O. Box 357 Hartshorne, OK 74547 TEL: (918) 297-7219

Pennsylvania Poultry Federation 500 North Progress Avenue Harrisonburg, PA 17109 TEL: (717) 652-7530 FAX: (717) 652-0230

South Carolina Poultry Federation 1201 Main Street, Suite 1220 AT&T Building Columbia, SC 29201 TEL: (803) 748-1283

FAX: (803) 748-1294

Tennessee Egg and Poultry Association P.O. Box 11082 Knoxville, TN 37939-1082 TEL: (615) 974-7351 FAX: (615) 974-7448

Tennessee Poultry Improvement Board P.O. Box 40627 Nashville, TN 37204 TEL: (615) 360-0120 FAX: (615) 781-5309



Texas Poultry Federation 8130 Burnet Road P.O. Box 9589 Austin, TX 78766-9589 TEL: (512) 451-6816 FAX: (512) 454-4221

APPLICATION AGREEMENT FORMS



LAND APPLICATION AGREEMENT FORMS

The following agreement forms will help producers with manure management liability issues. The forms will allow producers to form cooperative agreements with other parties for handling and disposal of manure. The forms were taken from those in the North Carolina "Dry Poultry Litter" Handbook.







Example 1 EXAMPLE SHARED RESPONSIBILITY AGREEMENT FOR LAND APPLICATION OF POULTRY LITTER

1.	anc p				
	owne	d by	ered by this agreement was remove	located at	
	*****		in		County
			be managed to ensure that there is		the surface waters.
	D.	Litter will not	be stockpiled within 100 feet of per g and providing a copy of the waste	ennial streams of wells.	the recognibility of the producer
	Ç.	information w	g and providing a copy of the wast ill be used in determining the applic and user as appropriate.	eation rate. The applicator w	ill provide the waste analysis to the
	d.	Application ra	tes will be calculated to not exceed	the agronomic use or phospl	horous needs whichever is more lin
	e.		pplied at rates not to exceed the agr		д стор.
	f.	Soils from fiel	ds receiving poultry litter will be sa	mpled annually.	
	g.	recommendati		l be maintained in accordance	ce with the standard soil test report
	h.		g for land application.		
			vill be kept for three years of soil te		
		Records v	ill be kept for three years indicating	the location of fields and re	ates at which litter is land annlied
II.	I the	landowner or la	nd user of fields where litter is appl from part	ied accept responsibility for	
II. —	I the	landowner or la	nd user of fields where litter is appl	ied accept responsibility for I.	
II. —	I the	landowner or la	nd user of fields where litter is appl from part	ied accept responsibility for I. Phone	items <u>a,b,h</u> and
111		landowner or la	nd user of fields where litter is appl from part Print Name	ied accept responsibility for I. Phone	items <u>a,b,h</u> ande Number
111	. I the	landowner or la	nd user of fields where litter is appl from part Print Name Signature	ied accept responsibility for I. Phone	items <u>a,b,h</u> ande Number
111	. I the	landowner or la	nd user of fields where litter is appl from part Print Name Signature	ied accept responsibility for I. Phone Phone The property of the property o	items <u>a,b,h</u> ande Number





${\tt Example~2} \\ {\tt SHARED~RESPONSIBILITY~AGREEMENT~FOR~LAND~APPLICATION~OF~POULTRY~LITTER} \\$

	owned by		from the poultry operation located at			
	inCounty.					
	 a. The litter must be managed to ensure that there is no discharge of the litter to the surface waters. b. Litter will not be stockpiled within 100 feet of perennial streams or wells. 					
	c. Waste san information landowne	npling and providing a copy of the waste analysis to on will be used in determining the application rate. T r'or land user as appropriate.	the applicator is the responsibility of the producer. This he applicator will provide the waste analysis to the			
	requireme	lication rates will be calculated and applied at rates nents whichever is more limiting of the receiving crop.				
		n fields receiving litter will be sampled annually, for all fields receiving litter will be maintained in accordations	ordance with the standard soil test report			
	g. Record ke 1) Soil t	seeping for land application. est and waste analysis will be kept for three years. I location and litter application records will be kept for	or three years.			
II.	I	*, representing	** accept responsibility for items			
	0:	from part I. Date	Phone Number			
	Signature	The second secon	The state of the s			
	I	*, representing	** accept responsibility for items			
	I	*, representing from part l.				
	ISignature	*, representing from part IDate , representing	** accept responsibility for items			
	ISignature	*, representingfrom part I. Date *, representingfrom part I.	** accept responsibility for items Phone Number			
	I	*, representingfrom part I	** accept responsibility for items Phone Number ** accept responsibility for items			



DRY LITTER UTILIZATION - THIRD PARTY APPLICATOR AGREEMENT (EXAMPLE)

i,	hereby acknowled	ge that I have received a Dry	Pitter Management Lig
that was developed for n	ny facility located at		
in	_ County.		
I hereby agree to manag manner to include the re records as specified in the	quirements of my Dry Litter	try litter that I received from Management Plan and keep o	the facility below in a copies of all required
Third Party Receiver:		Date:	
Manure Producer:		Date:	
Terms of Agreement:	, 19	to,	·





REFERENCES

- Arkansas Soil and Water Conservation Commission and the Water Resources Center, No Date, Composting: A safe and simple alternative in water quality and poultry disposal pits. Fact Sheet 2, Arkansas Soil and Water Conservation Commission, Little Rock.
- Barker, J.C., 1990, Livestock waste sampling, analysis, and calculation of land application rates. Reprint. EBAE 111-84. Agricultural Extension Service, North Carolina State University, Raleigh.
- Bell, D., 1982, Marketing poultry manure. University of California, Riverside.
- Blake, J.P., 1993, Mortality Management Presentation; Poultry Waste Management and Water Quality Workshop. Southeastern Poultry and Egg Association, Atlanta, GA.
- Brandel, V.A., 1988, Soil phosphorous,: managing it effectively, Fact Sheet 513. Cooperative Extension Service, University of Maryland, College Park.
- Brodie, H.L. and V.A. Bradel, 1990, Manure testing, FS-430. Cooperative Extension Service, University of Maryland, College Park.
- Brodie, H.L., L.E. Carr, and C.F. Miller, 1990, Structures for broiler litter manure storage, Fact Sheet 416, Cooperative Extension Service, University of Maryland, College Park.
- Brown, W.R., 1993, Composting poultry manure. Presentation; Poultry Waste Management and Water Quality Workshop, Southeastern Poultry and Egg Association, Atlanta, GA.
- Cabe Associates, Inc., 1991, Poultry manure storage and process: alternative evaluation. Final Report. Project 100-286. University of Delaware Research and Education Center, Georgetown.
- Carter, T.A., and R.E. Sneed, 1987, Drinking water quality for poultry. PS&T Guide No. 42. Cooperative Extension Service, North Carolina State University, Raleigh.
- Department of Natural Resources and Environmental Control, 1989, Poultry manure management: a supplement to Delaware guidelines. Cooperative Bulletin 24, Delaware Cooperative Extension, University of Delaware, Newark.



- Donald, J.O. and J.P. Blake, 1990, Economics of transporting poultry litter as a fertilizer, DTP Circular 10/90-007. Alabama Cooperative Extension Service. , 1990, Use and construction of poultry burial pits, DTP Circular 10/90-013, Alabama Cooperative Extension Service, Auburn University, Auburn, AL. 1990, Installation and use of incincrators, DPT Circular 11/90-014. Alabama Cooperative Extension Service, Auburn University, Auburn, AL. Fulhage, C., 1989, Reduce environmental problems with proper land application of animal wastes. WQ201. University Extension, University of Missouri, Columbia. 1990, Reduce environmental problems with proper land application of animal waste. WQ201. Cooperative Extension Service, University of Missouri, Columbia. 1992, Reduce environmental problems with proper land application of animal wastes, WQ201, University Extension, University of Missouri, Columbia. 1992, Composting poultry carcasses in Missouri, WQ205. Cooperative Extension, University of Missouri, Columbia. 1992, Land application considerations for animal wastes. WQ202, University Extension, University of Missouri, Columbia. 1992, National Engineering Handbook 210, Part 651, In Agricultural Waste Management Handbook, Soil Conservation Services, Washington, DC. Goan, H.C. and J. Jared, 1991, Poultry manure--proper handling and application to protect our water resources. PB 1421. Cooperative Extension Service, University of Tennessee, Knoxville.
- Keeney, D.R. and R.F. Follett, 1991, Overview and introduction. Chapter 1, <u>In</u>

 Managing nitrogen for groundwater quality and farm profitably. Proceedings.

 Soil Science Society of America, Madison, WI.
- Killpack, S. and D. Bucholz, 1991, What is nitrogen? WQ2251. University Extension, University of Missouri, Columbia.
- Margette, W.L. and R.A. Weismiller, 1991, Nutrient management for waste quality protection, Bay Fact Sheet 3, Cooperative Extension Service, University of Maryland, College Park.



- Missouri Department of Natural Resources, 1993, Obtaining a DNR letter of approval or a livestock waste management system, WQ217, University Extension, University of Missouri, Columbia.
- Murphy, D.W. and Carr, L.E., 1991, rev. Composting Dead Birds. Fact Shect 537, Cooperative Extension, University of Maryland, College Park.
- Ogburn, C.B. and J.O. Donald, 1990, Calibrating spreaders for the application of animal and poultry manure, DPT Circular 11/90-006. Cooperative Extension Service, Auburn University, Auburn, AL.
- Poultry Water Quality Handbook, 1994, Poultry Water Quality Consortium, HB-2C, 1101 Market Street, Chattanooga, TN 37402-2801.
- Swanson M.H., No Date, Some reflections on dried poultry waste, University of California Cooperative Extension, Riverside.
- U.S. Department of Agriculture, 1991, Improving and protecting water quality in Georgia... one drop at a time, Soil Conservation Service, Athens, GA.
- , 1992, National Engineering Handbook 210, Part 651. In Agriculture Waste Management Field Handbook, Soil Conservation Service, Washington DC.
- U.S. Environmental Protection Agency, 1993, The watershed protection approach: a project focus, draft, Assessment and Watershed Protection Division. Washington, DC.
- Wells, K.L., G.W. Thomas, J.L. Sims, and M.S. Smith, 191, Managing soil nitrates for agronomic efficiency and environmental protection. AGR-147. Cooperative Extension Service, University of Kentucky, Lexington.
- Wineland, M.J. and Carter, T.A., 1987, Poultry science and technology guide--proper disposal of dead poultry. PS&T Guide No. 19, North Carolina Agricultural Extension Service, Raleigh.
- Woodward, M., No Date, Manure spreader calibration worksheet. Technical Note #4, Cooperative Extension Service, Pennsylvania State University, Lancaster.
- Zimmerman, M., 1993, Freezing for the rendering plant, Presentation, Poultry Waste Management and Water Quality Workshop. Southeastern Poultry and Egg Association, Atlanta, GA.
- Zublena, J., 1993, Land applications: nutrient management plan. Presentation, Poultry Waste Management and Water Quality Workshop, Southeastern Poultry and Egg Association, Atlanta, GA.





Zublena, J.P., Barker, J.C., and Carter, T.A., 1993, Soil facts: poultry manure as a fertilizer source. AG-439-5. WQWM-41. North Carolina Cooperative Extension Service, Raleigh.

